

# 3-enumerated alternating sign matrices

Yu. G. Stroganov

*Institute for High Energy Physics  
142284 Protvino, Moscow region, Russia*

## Abstract

Let  $A(n, r; 3)$  be the total weight of the alternating sign matrices of order  $n$  whose sole '1' of the first row is at the  $r^{th}$  column and the weight of an individual matrix is  $3^k$  if it has  $k$  entries equal to -1. Define the sequence of the generating functions  $G_n(t) = \sum_{r=1}^n A(n, r; 3)t^{r-1}$ .

Results of two different kind are obtained. On the one hand I made the explicit expression for the even subsequence  $G_{2\nu}(t)$  in terms of two linear homogeneous second order recurrence in  $\nu$  (Theorem 1). On the other hand I brought to light the nice connection between the neighbouring functions  $G_{2\nu+1}(t)$  and  $G_{2\nu}(t)$  (Theorem 2).

The 3-enumeration  $A(n; 3) \equiv G_n(1)$  which was found by Kuperberg is reproduced as well.


## 1 Introduction and results

An alternating sign matrix, or ASM, is a matrix of 0's, 1's, and -1's such that the non-zero elements in each row and column alternate between 1 and -1 and begin and end with 1. Recall the conventional notations:

- $A(n)$  the number of  $n \times n$  ASMs;
- $A(n; 3)$  the total weight of all  $n \times n$  ASMs, where the weight of an individual matrix is  $3^k$  if it has  $k$  entries equal to -1;
- $A(n, r)$  the number  $n \times n$  ASMs for which the (unique) '1' of the first row is at the  $r^{th}$  column;

- $A(n, r; 3)$  the total weight of  $n \times n$  ASMs for which the (unique) ‘1’ of the first row is at the  $r^{th}$  column and the weight of an individual matrix is  $3^k$  if it has  $k$  entries equal to -1;

An ultra-short background can be described by the scheme:



We investigate the generating function  $G_n(t) = \sum_{r=1}^n A(n, r; 3) t^{r-1}$ . The next results follow directly from the general method elaborated in paper [5].

**Theorem 1** (ASMs of the even order) The normalized generating function  $\tilde{G}_{2\nu+2}(t) \equiv G_{2\nu+2}(t)/A(2\nu+2; 3)$  is given by a formula

$$\tilde{G}_{2\nu+2}(t) = \frac{(2\nu+1)!}{(3\nu+2)!} \frac{\{(3\nu+2)g_\nu^{(1)}(t) - (3\nu+1)g_\nu^{(2)}(t)\}}{t+1}, \quad \nu \geq 0$$

where  $g_\nu^{(j)}(t)$ ,  $(j = 1, 2)$  are two sequences of the polynomials which are fixed by the second order recurrence

$$\begin{aligned} & 3(1-t^2)^2 g_{\nu+1}^{(j)}(t) - \\ & -(2\nu+1)(1+4t+t^2)[3(1+t+t^2)^2 - (1+4t+t^2)^2] g_\nu^{(j)}(t) - \\ & -(9\nu^2 - j^2)t^2(1+2t)^2(2+t)^2 g_{\nu-1}^{(j)}(t) = 0, \quad (j = 1, 2) \end{aligned}$$

with the initial data

$$\begin{aligned} g_0^{(1)}(t) &= (t^2 + t + 1)/3, \quad g_1^{(1)}(t) = [(1+2t)^2(2+t)^2 - 9t^2]/18 \\ g_0^{(2)}(t) &= -(t^2 + 4t + 1)/3, \quad g_1^{(2)}(t) = -[(1+2t)^2(2+t)^2 + 9t^2]/18. \end{aligned}$$

**Theorem 2** The normalized generating functions  $\tilde{G}_n(t) \equiv G_n(t)/A(n; 3)$  satisfy

$$\tilde{G}_{2\nu+1}(t) = \tilde{G}_{2\nu}(t) \frac{2(1+2t)(2+t)}{9(t+1)}, \quad \nu > 0$$

As far as the normalizing factor  $A(n; 3)$  is concerned one can easily reproduce Theorem 2 of Kuperberg paper [4] using equations of Section 4.

## 2 Outline of method

This small note can be considered as the supplement to my recent paper [5]. I use the notation and results of the cited paper without reviewing them.

Consider the inhomogeneous six-vertex model with the domain wall boundary conditions [6]. Spectral parameters  $\{x_1, x_2, \dots, x_n\}$  and  $\{y_1, y_2, \dots, y_n\}$  are attached to the horizontal and vertical lines respectively. We fix them as follows:

$$\begin{aligned} x_1 &= u, \\ x_i &= \pi/2, \quad i = 2, 3, \dots, n, \\ y_i &= 0, \quad i = 1, 2, \dots, n. \end{aligned}$$

The 3-enumeration of ASMs corresponds to the 6-vertex model with the special crossing parameter  $\eta = 2\pi/3$ , so that the Boltzmann weights in the upper row are given by:

$$\begin{aligned} a(u) &= 2 \sin(u + \pi/3), \quad b(u) = 2 \sin(u - \pi/3), \\ c(u) &= \sqrt{3}. \end{aligned} \tag{1}$$

All weights in the remaining rows<sup>1</sup> correspond to the 3-enumeration:

$$a = b = 1 \quad c = \sqrt{3}.$$

The partition function of the model is given by the trigonometric polynomial of degree  $n - 1$ :

$$Z(u) = \sqrt{3} \sum_{r=1}^n A(n, r; 3) a^{r-1}(u) b^{n-r}(u). \tag{2}$$

---

<sup>1</sup>We use formulae (1) of paper [5] changing the normalization by a factor  $\sqrt{3}$ .

Using equations (1), (2) and right-left symmetry of ASMs we obtain that  $Z(u)$  satisfy

$$Z(-u) = (-1)^{n-1} Z(u). \quad (3)$$

It was shown in paper [5] (see equations (5-7) of the cited paper) that one has the equation

$$f(u) + f(u + \frac{2\pi}{3}) + f(u + \frac{4\pi}{3}) = 0 \quad (4)$$

with

$$f(u) = Z(u) \sin^n(u) \cos^{n-1}(u). \quad (5)$$

We see that the function  $f(u)$  is an odd trigonometric polynomial of degree  $3n - 2$ . It turns out that under the circumstances the function  $f(u)$  and consequently the partition function  $Z(u)$  can be found up to an arbitrary constant. Solving these equations (the solution is given in Appendix A) we obtain

$$\begin{aligned} f_{2\nu+2}(u) = c_\nu & \left\{ (3\nu+2) \sum_{\alpha=0}^{\nu} \binom{\nu - \frac{1}{3}}{\alpha} \binom{\nu + \frac{1}{3}}{\nu - \alpha} \sin(2 - 6\nu + 12\alpha)u - \right. \\ & \left. - (3\nu+1) \sum_{\alpha=0}^{\nu} \binom{\nu - \frac{2}{3}}{\alpha} \binom{\nu + \frac{2}{3}}{\nu - \alpha} \sin(4 - 6\nu + 12\alpha)u \right\} \end{aligned} \quad (6)$$

for  $n$  even and

$$f_{2\nu+1}(u) = r_\nu \cos 3u f_{2\nu}(u) \quad (7)$$

for  $n$  odd, where multipliers  $c_\nu$  and  $r_\nu$  do not depend on  $u$ .  $\nu$  equals to the integer part of  $(n-1)/2$ .

Formulae (1), (2), (5), (6) and (7) solve in a sense the problem. We are left with the task of elaborating an effective method for explicit calculation of  $A(n, r; 3)$ . Besides we have to find the multipliers  $c_\nu$  and  $r_\nu$ .

### 3 Recursion relations

Let us begin with  $n$  even. It is convenient to present equation (6) as follows

$$f_{2\nu+2}(u) = c_\nu [(3\nu+2) F_\nu^{(1)}(u) - (3\nu+1) F_\nu^{(2)}(u)] \quad (8)$$

where  $F_\nu^{(j)}(u)$ ,  $(j = 1, 2)$  are trigonometric polynomials

$$F_\nu^{(j)}(u) = \sum_{\alpha=0}^{\nu} \binom{\nu - \frac{j}{3}}{\alpha} \binom{\nu + \frac{j}{3}}{\nu - \alpha} \sin 2(j - 3\nu + 6\alpha)u$$

One can verify directly that these polynomials satisfy a recurrence

$$9\nu(\nu + 1)F_{\nu+1}^{(j)}(u) - 18\nu(2\nu + 1)\cos 6u F_\nu^{(j)}(u) - 4(9\nu^2 - j^2)\sin^2 6u F_{\nu-1}^{(j)}(u) = 0, \quad (j = 1, 2) \quad (9)$$

with the initial data

$$\begin{aligned} F_0^{(1)}(u) &= \sin 2u, & F_1^{(1)}(u) &= \frac{2}{3}(\sin 8u - 2\sin 4u) \\ F_0^{(2)}(u) &= \sin 2u, & F_1^{(2)}(u) &= \frac{1}{3}(\sin 10u - 5\sin 2u) \end{aligned}$$

It was shown in paper [7] that quotients

$$\Phi_\nu^{(j)}(w) = F_\nu^{(j)}(u) / \sin^{2\nu+1} 2u, \quad (j = 1, 2) \quad (10)$$

are polynomials in the variable  $w = \cos 2u$ . Recurrence (9) can be rewritten as a recurrence for these polynomials:

$$9\nu(\nu + 1)(1 - w^2)\Phi_{\nu+1}^{(j)}(w) + 18\nu(2\nu + 1)w(3 - 4w^2)\Phi_\nu^{(j)}(w) - 4(9\nu^2 - j^2)(1 - 4w^2)^2\Phi_{\nu-1}^{(j)}(w) = 0, \quad (j = 1, 2) \quad (11)$$

with the initial data

$$\begin{aligned} \Phi_0^{(j)}(w; 1) &= 1, & \Phi_1^{(j)}(w; 1) &= -16w/3 \\ \Phi_0^{(j)}(w; 2) &= 2w, & \Phi_1^{(j)}(w; 2) &= -4(1 + 4w^2)/3 \end{aligned} \quad (12)$$

Combining equations (2), (5), (8) and (10) we obtain firstly

$$\begin{aligned} Z(u) \sin u &\equiv \sqrt{3} \sin u a^{2\nu+1}(u) \sum_{r=1}^{2\nu+2} A(2\nu + 2, r; 3) \left( \frac{b(u)}{a(u)} \right)^{2\nu+2-r} = \\ &= 2^{2\nu} c_\nu \{ (3\nu + 2) \Phi^{(1)}(w) - (3\nu + 1) \Phi^{(2)}(w) \}, \end{aligned}$$

and using a new variable  $t$

$$t = \frac{b(u)}{a(u)} = \frac{\sin(u - \pi/3)}{\sin(u + \pi/3)}, \quad (13)$$

we can present the last equation as follows

$$\begin{aligned}
G_{2\nu+2}(t) &\equiv \sum_{r=1}^{2\nu+2} A(2\nu+2, r; 3) t^{2\nu+2-r} = \\
&= \left(\frac{4}{3}\right)^{\nu+1} \frac{c_\nu}{\sqrt{3}(t+1)} (t^2+t+1)^{\nu+1} \\
&\quad \left\{ (3\nu+2) \Phi_\nu^{(1)}\left(-\frac{t^2+4t+1}{2(t^2+t+1)}\right) - (3\nu+1) \Phi_\nu^{(2)}\left(-\frac{t^2+4t+1}{2(t^2+t+1)}\right) \right\}
\end{aligned} \tag{14}$$

where  $c_\nu$  is a multiplier which does not depend on  $t$  (see equation (6)).

It is clear that  $A(2\nu+2, r; 3) = A(2\nu+2, 2\nu+3-r; 3)$  therefore we can write

$$G_{2\nu+2}(t) \equiv \sum_{r=1}^{2\nu+2} A(2\nu+2, r; 3) t^{r-1}$$

in the first part of equation (14).

On the other hand combining equations (2), (5) and (7) we obtain firstly the relation between the partition functions:

$$Z_{2\nu+1}(u) \sin u \cos u = r_\nu \cos 3u Z_{2\nu}(u),$$

and then transform it into the equation connecting the generating functions

$$G_{2\nu+1}\left(\frac{b(u)}{a(u)}\right) a(u) \sin u = r_\nu \frac{\cos 3u}{\cos u} G_{2\nu}\left(\frac{b(u)}{a(u)}\right).$$

Using the variable  $t$  (see equation (13)) we rewrite the last equation as

$$G_{2\nu+1}(t) = -r_\nu \frac{2(1+2t)(2+t)}{3(1+t)} G_{2\nu}(t) \tag{15}$$

where  $r_\nu$  is a multiplier which does not depend on  $t$ .

## 4 Multipliers $c_\nu$ and $r_\nu$

We are prepared now to calculate the multipliers  $c_\nu$  and  $r_\nu$ . The point that for some special values of the spectral parameter  $u$  (see equation (1)) the recurrences we consider are simplified and can be easily solved. Details can be found in Appendix B. Let us compare equations (30), (31), (34) and (35). Excluding the multipliers from these equations we obtain

$$\frac{A(2\nu + 2; 3)}{A(2\nu + 1; 3)} = \frac{3^\nu \nu! (3\nu + 2)!}{[(2\nu + 1)!]^2}$$

and

$$\frac{A(2\nu + 1; 3)}{A(2\nu; 3)} = \frac{9A(2\nu; 3)}{4A(2\nu - 1; 3)} = \frac{3^\nu \nu! (3\nu)!}{[(2\nu)!]^2}.$$

Using these fractions and the trivial equation  $A(1; 3) = 1$  one can reproduce Theorem 2 of Kuperberg paper [4]. Fixing the multipliers  $c_\nu$  from equation (34), inserting them into (14) and defining new polynomials:

$$g_\nu^{(j)}(t) = \frac{\nu! (t^2 + t + 1)^{\nu+1}}{3 \cdot 4^\nu} \Phi_\nu^{(j)} \left( -\frac{t^2 + 4t + 1}{2(t^2 + t + 1)} \right)$$

we obtain our Theorem 1. On the other hand we easily obtain the Theorem 2 normalizing the generation function entering equation (15).

*Acknowledgments* The work was supported in part by the Russian Foundation for Basic Research under grant # 01-01-00201 and by the INTAS under grant # 00-00561.

### Appendix A. Calculation of $f(u)$ up to arbitrary constant

Recall that  $f(u)$  is an odd trigonometric polynomial of degree  $3n - 2$ . We can therefore write

$$f(u) = b_1 \sin(3n - 2)u + b_2 \sin(3n - 4)u + b_3 \sin(3n - 6)u + \dots \quad (16)$$

Equation (4) is satisfied if and only if  $b_{3\kappa} = 0$ . This condition implies

$$f(u) = \sum_{k=0}^{n-1} \beta_k \sin(4 - 3n + 6k)u. \quad (17)$$

According to equation (5) the monomial  $\sin^n u \cos^{n-1} u$  divides this function and we obtain a set of equations

$$\begin{aligned} f^{(m)}(u)|_{u=0} &= 0 \quad m = 0, 1, \dots, n-1, \\ f^{(m)}(u)|_{u=\pi/2} &= 0 \quad m = 0, 1, \dots, n-2, \end{aligned} \quad (18)$$

where  $f^{(m)}$  the derivative of  $f$  of the order  $m$ . Let us insert equation (17) into (18).

If  $n$  is even then we obtain the system:

$$\begin{aligned} \sum_{k=0}^{2\nu+1} \beta_k (3k - 1 - 3\nu)^{2\mu+1} &= 0, \quad \mu = 0, 1, \dots, \nu \\ \sum_{k=0}^{2\nu+1} (-1)^k \beta_k (3k - 1 - 3\nu)^{2\mu+1} &= 0, \quad \mu = 0, 1, \dots, \nu - 1. \end{aligned}$$

Considering separately odd and even  $k$  and combining obtained relations we get two similar systems

$$\sum_{\kappa=0}^{\nu} \beta_{2\kappa} (6\kappa - 1 - 3\nu)^{2\mu+1} = 0, \quad \mu = 0, 1, \dots, \nu - 1, \quad (19)$$

$$\sum_{\kappa=0}^{\nu} \beta_{2\kappa+1} (6\kappa + 2 - 3\nu)^{2\mu+1} = 0, \quad \mu = 0, 1, \dots, \nu - 1, \quad (20)$$

and one additional relation:

$$\sum_{\kappa=0}^{\nu} \beta_{2\kappa} (6\kappa - 1 - 3\nu)^{2\nu+1} + \sum_{\kappa=0}^{\nu} \beta_{2\kappa+1} (6\kappa + 2 - 3\nu)^{2\nu+1} = 0. \quad (21)$$

Systems (19) and (20) are equivalent to the condition that the relations

$$\sum_{\kappa=0}^{\nu} \beta_{2\kappa} (6\kappa - 1 - 3\nu) p((6\kappa - 1 - 3\nu)^2) = 0, \quad (22)$$

and

$$\sum_{\kappa=0}^{\nu} \beta_{2\kappa+1} (6\kappa + 2 - 3\nu) p((6\kappa + 2 - 3\nu)^2) = 0, \quad (23)$$

are valid for all polynomials  $p(x)$  of degree  $\nu - 1$ . Let us begin with system (22). Consider  $\nu$  special polynomials of degree  $\nu - 1$ :

$$p_{\alpha}(x) = \prod_{\kappa=1, \kappa \neq \alpha}^{\nu} (x - (6\kappa - 1 - 3\nu)^2) \quad \alpha = 1, 2, \dots, \nu.$$

Inserting these polynomials into (22) we find simple relations connecting  $\beta_{2\alpha}$  with  $\beta_0$  which can be written as

$$\beta_{2\alpha} = c_1 \binom{\nu + \frac{1}{3}}{\alpha} \binom{\nu - \frac{1}{3}}{\nu - \alpha} \quad \alpha = 0, \dots, \nu. \quad (24)$$

In the same way we convert equation (23) into

$$\beta_{2\alpha+1} = c_2 \binom{\nu - \frac{2}{3}}{\alpha} \binom{\nu + \frac{2}{3}}{\nu - \alpha} \quad \alpha = 0, \dots, \nu. \quad (25)$$

We have to satisfy the additional relation (21). Taking into account equations (24) and (25) one can reduce this relation to the equation:

$$c_2/c_1 = - \frac{\sum_{\alpha=0}^{\nu} \binom{\nu + \frac{1}{3}}{\alpha} \binom{\nu - \frac{1}{3}}{\nu - \alpha} (6\alpha - 1 - 3\nu)^{2\nu+1}}{\sum_{\alpha=0}^{\nu} \binom{\nu - \frac{2}{3}}{\alpha} \binom{\nu + \frac{2}{3}}{\nu - \alpha} (6\alpha + 2 - 3\nu)^{2\mu+1}}.$$

Using Mathematica I found that the right side fraction probably equals to  $(3\nu + 1)/(3\nu + 2)$ . Then I appealed to the domino forum and a few days a proof of more general identity was given by Guoce Xin [8]. A simpler proof of Guoce Xin's identity was presented by Ira Gessel [9].

As a result we obtain (up to an arbitrary constant multiplier):

$$\begin{aligned} f(u) &\equiv f_{2\nu+2}(u) \propto \\ &\sim \left\{ (3\nu + 2) \sum_{\alpha=0}^{\nu} \binom{\nu + \frac{1}{3}}{\alpha} \binom{\nu - \frac{1}{3}}{\nu - \alpha} \sin(-2 - 6\nu + 12\alpha)u + \right. \\ &\quad \left. + (3\nu + 1) \sum_{\alpha=0}^{\nu} \binom{\nu - \frac{2}{3}}{\alpha} \binom{\nu + \frac{2}{3}}{\nu - \alpha} \sin(4 - 6\nu + 12\alpha)u \right\}. \end{aligned} \quad (26)$$

Changing  $\alpha$  with  $\nu - \alpha$  in the first sum we obtain a variation of this formula which is used in the body of the paper.

If  $n$  is odd then inserting (17) into (18) we obtain

$$\begin{aligned} \sum_{k=0}^{2\nu} \beta_k (6k + 1 - 6\nu)^{2\mu+1} &= 0, \quad \sum_{k=0}^{2\nu} (-1)^k \beta_k (6k + 1 - 6\nu)^{2\mu} = 0, \\ \mu &= 0, 1, \dots, \nu - 1, \end{aligned}$$

where  $n = 2\nu + 1$ . Consider separately  $k$  odd and even:

$$\begin{aligned} \sum_{\kappa=0}^{\nu} \beta_{2\kappa} (12\kappa + 1 - 6\nu)^{2\mu+1} + \sum_{\kappa=1}^{\nu} \beta_{2\kappa+1} (12\kappa - 5 - 6\nu)^{2\mu+1} &= 0, \\ \sum_{\kappa=0}^{\nu} \beta_{2\kappa} (12\kappa + 1 - 6\nu)^{2\mu} - \sum_{\kappa=1}^{\nu} \beta_{2\kappa+1} (12\kappa - 5 - 6\nu)^{2\mu} &= 0, \\ \mu &= 0, 1, \dots, \nu - 1. \end{aligned}$$

This system is equivalent to the condition that the relation

$$\sum_{\kappa=0}^{\nu} \beta_{2\kappa} p(12\kappa + 1 - 6\nu) - \sum_{\kappa=1}^{\nu} \beta_{2\kappa+1} p(-12\kappa + 5 + 6\nu) = 0 \quad (27)$$

is valid for all polynomials  $p(x)$  of degree  $2\nu - 1$ . Consider  $2\nu$  special polynomials of degree  $2\nu - 1$ :

$$\begin{aligned} p_{\alpha}(x) &= \prod_{\kappa=1, \kappa \neq \alpha}^{\nu} (x - 1 + 6\nu - 12\kappa) \prod_{\kappa=1}^{\nu} (x - 5 - 6\nu + 12\kappa) \\ \tilde{p}_{\alpha}(x) &= \prod_{\kappa=1}^{\nu} (x - 1 + 6\nu - 12\kappa) \prod_{\kappa=1, \kappa \neq \alpha}^{\nu} (x - 5 - 6\nu + 12\kappa) \\ \alpha &= 1, 2, \dots, \nu. \end{aligned}$$

Inserting these polynomials into (27) we find simple relations connecting  $\beta_{\alpha}$  with  $\beta_0$ . It is possible to write the answer in terms of binomial coefficients:

$$\begin{aligned} f(u) &\equiv f_{2\nu+1} \propto \\ &\sim \left\{ \sum_{\alpha=0}^{\nu} \binom{\nu - \frac{2}{3}}{\alpha} \binom{\nu - \frac{1}{3}}{\nu - \alpha} \sin(1 - 6\nu + 12\alpha)u - \right. \\ &\quad \left. - \sum_{\alpha=0}^{\nu-1} \binom{\nu - \frac{1}{3}}{\alpha} \binom{\nu - \frac{2}{3}}{\nu - \alpha - 1} \sin(5 - 6\nu + 12\alpha)u \right\}. \end{aligned}$$

One can check directly that multiplying  $f_{2\nu+2}$  (see equation (26)) by  $\cos 3u$  one obtain  $f_{2\nu+3}$  which is given by the last equation with  $\nu \rightarrow \nu + 1$ .

## Appendix B. Generating functions in special points

Let us begin with the point  $u = \pi/3$ . For this case variable  $t$  which is defined by equation (13) equals to zero and the generating function  $G_n(t)$  comes to the first term  $A(n, 1; 3)$ . According to the trivial bijection between

the ASMs of order  $n$  whose sole '1' of the first row is at the first column and all ASMs of order  $n - 1$  one can write  $A(n - 1; 3)$  instead of  $A(n, 1; 3)$ . Equation (11) reduces to ( $w \equiv \cos 2u = -1/2$ )

$$3(\nu + 1) \Phi_{\nu+1}^{(j)}(-1/2) - 8(2\nu + 1) \Phi_{\nu}^{(j)}(-1/2) = 0, \quad (j = 1, 2) \quad (28)$$

According to equation (12) we have the initial data

$$\Phi_{\nu+1}^{(j)}(-1/2) = (-1)^{j+1} \frac{8}{3}, \quad (j = 1, 2) \quad (29)$$

Recurrence (28) with initial data (29) gives

$$\Phi_{\nu+1}^{(j)}(-1/2) = (-1)^{j+1} \left(\frac{4}{3}\right)^{\nu} \frac{(2\nu)!}{(\nu!)^2}.$$

and equation (14) comes to

$$A(2\nu + 1; 3) = \left(\frac{4}{3}\right)^{2\nu+1} c_{\nu} \frac{\sqrt{3} (2\nu + 1)!}{(\nu!)^2}. \quad (30)$$

It is clear that equation (15) reduces to

$$A(2\nu; 3) = -\frac{4}{3} r_{\nu} A(2\nu - 1; 3). \quad (31)$$

Consider now another convenient value of the spectral parameter  $u = \pi/2$ . Unlike the previous case variable  $t$  which is defined by equation (13) equals to unity and the generating function  $G_n(t)$  come to 3-enumeration of all ASMs of order  $n$ :

$$G_n(1) = A(n; 3).$$

Equation (11) reduces to ( $w \equiv \cos 2u = -1$ )

$$\nu (2\nu + 1) \Phi_{\nu}^{(j)}(-1) - 2 (9 \nu^2 - j^2) \Phi_{\nu-1}^{(j)}(-1) = 0, \quad (j = 1, 2) \quad (32)$$

with the initial data

$$\begin{aligned} \Phi_0^{(1)}(-1) &= 1, & \Phi_1^{(1)}(-1) &= 16/3 \\ \Phi_0^{(2)}(-1) &= -2, & \Phi_1^{(2)}(-1) &= -20/3. \end{aligned} \quad (33)$$

Recurrence (32) with initial data (33) gives

$$\begin{aligned}\Phi_{\nu}^{(1)}(-1) &= \left(\frac{4}{3}\right)^{\nu} \frac{(3\nu+1)!}{\nu! (2\nu+1)!} \\ \Phi_{\nu}^{(2)}(-1) &= -\left(\frac{4}{3}\right)^{\nu} \frac{(3\nu+2)(3\nu)!}{\nu! (2\nu+1)!}.\end{aligned}$$

and equation (14) comes to

$$A(2\nu+2; 3) = \left(\frac{16}{3}\right)^{\nu} \frac{4 c_{\nu} (3\nu+2)!}{\sqrt{3} \nu! (2\nu+1)!}. \quad (34)$$

On the other hand equation (15) reduces to

$$A(2\nu+1; 3) = -3r_{\nu}A(2\nu; 3). \quad (35)$$

## References

- [1] W. H. Mills, D. P. Robbins, and H. Rumsey, *Proof of the Macdonald conjecture*, Invent. Math., **66** (1982) 73-87.  
W. H. Mills, D. P. Robbins, and H. Rumsey, *Alternating-sign matrices and descending plane partitions*, J. Combin. Theory Ser. A, **34** (1983) 340-359.
- [2] D. Zeilberger, *Proof of the alternating sign matrix conjecture*, Elec. J. Comb., **3(2)** (1996) R13.
- [3] D. Zeilberger, *Proof of the refined alternating sign matrix conjecture*, New York J. Math., **2** (1996) 59-68.
- [4] G. Kuperberg, *Another proof of the alternating-sign matrix conjecture*, Internat. Math. Res. Notes, **3** (1996) 139-150.
- [5] Yu. G. Stroganov, *A new way to deal with Izergin-Korepin determinant at root of unity*, arXiv:math-ph/0204042
- [6] V. E. Korepin, *Calculation of norms of Bethe wave functions*, Comm. Math. Phys. **86** (1982) 391-418.

- [7] Yu. G. Stroganov, *The importance of being odd*, J. Phys. A **34** (2001) L179-L185
- [8] G. Xin, Domino Forum, 12/24/2002
- [9] I. Gessel, Domino Forum, 01/10/2003